DR. JAMES C. SALATO WILDLIFE EDUCATION CENTER NATIVE PLANT PROGRAM

2010 Fall Sale Saturday, August 28, 2010

Kentucky Department of Fish and Wildlife Resources #1 Sportsman's Lane Frankfort KY 40601 (502) 564-7863 or (800) 858-1549

All Plants \$5.00 and \$6.00

NATIVE PERENNIALS

| _Agave (Agave virginiana) | 3-4' | succulent leaves, green flowers in summer, drought tolerant, full sun |
|--|-------|--|
| _Alumroot, Hairy (Heuchera villosa)) | 10" | small green-white flowers, soft mounds of foliage, good ground cover, shade, well-drained soil |
| _Aster, Aromatic (Aster oblongifolius) | 2-3' | blue flowers with yellow centers in fall, full or partial sun, drought tolerant |
| _Aster, Arrow-leaved (Aster sagittiflorus) | 3' | blue flowers, fall, large arrow-shaped leaves, partial shade, moist soil |
| _Aster, Calico (Aster laterifolius) | 3-4' | small white flowers in autumn, full sun, average soil |
| _Aster, Crooked-stemmed (Aster prenanthoides) | 1-2' | wide lavender flowers in autumn, full to partial sun, moist to wet soil |
| _Aster, New England (Aster novae-angliae) | 3-5' | large showy purple flowers, late bloomer, full sun, well-drained soil |
| _Aster, Short's (Aster shortii) | 2-3' | deep violet-blue flowers in fall, shade to partial shade, well-drained soil |
| _Aster, Smooth (Aster laevis) | 2-4' | blue, blue-gray foliage, fall, sun to partial shade, very drought tolerant, birds, butterflies, hummers |
| _Aster, Swamp (Aster puniceus) | 6' | large mass of violet flowers in Aug-Sept, sun to partial sun, wet to normal soil |
| _Aster, Western Silky (Aster sericeus) | 12" | pink-purple on slender stems, fall, soft silvery leaves, well-drained soil, rock gardens |
| _Aster, Willow (Aster praealtus) | 3-5' | lavender, late summer-fall, full sun, moist soil, wet prairies |
| Beardtongue, Small's (Penstemon smallii) | 1-2' | pink, early summer, dense spikes, avg. to rich, well-drained soil, sun to partial, birds, butterflies |
| Beardtongue, Smooth (Penstemon digitalis) | 3-4' | white, elegant plant, large basal leaves, early source of nectar, well-drained soil, sun to partial sun 🛪 🕷 |
| _Bee-Balm (Monarda didyma) | 3-4' | bright red flowers in dense clusters, mint, grows in avg. or moist soil, butterflies, hummers |
| _Bergamot, Wild (Monarda fistulosa) | 3-5' | lavender flowers, June, drought tolerant, partial sun, hummingbird, moths |
| _Blazing Star, Cylindrical (Liatris cylindracea) | 1-2' | pink feathery flowers, dainty, July-Sept., avg. to dry, well-drained soil, drought tolerant, rock garden, full sun |
| _Blazing Star, Prairie (Liatris pycnostachya) | 3-4' | dense spikes on erect stems, avg. to moist soil, full sun, butterflies, hummingbirds |
| _Blazing Star, Rough (Liatris aspera) | 3-5' | pink-purple button-shaped flower clusters, Aug-Oct, full sun, average to dry soil, hummers, butterflies |
| _Blue Sage (Salvia azurea) | 3-5' | blue 2-lipped flowers, July, well-drained soil, hummingbirds, butterflies |
| _Blazing Star, Spiked (Liatris spicata) | 4-6' | pink dense spikes on erect stems, avg-well-drained soil, drought tolerant and adapted to clay soils, full sun |
| _Blue-eyed grass (Sisyrinchium angustifolium) | 6-18" | iris family with grass-like leaves, blue flowers, May-July, well drained soil, sun to partial shade |
| _Blue Star (Amsonia tabernaemontana) | 3-4' | blue star-shaped flowers, late spring, full or partial sun, moist soil |
| _Boltonia (Boltonia asteroides) | 2-6' | white disk flowers, Sep-Oct, tolerates wide range of soil and moisture |
| | | |

| Boneset (Eupatorium perfoliatum) | 3-5' | white, August, sun to partial sun, wet ground, butterflies |
|--|-------|---|
| Bradbury's Monarda (Monarda bradburnia) | 2-3 | light pink blooms, May, full to partial sun, drought-tolerant, hummingbirds, moths ** ** |
| Cactus, Prickly Pear (Opuntia humifusa) | 2' | showy yellow flowers, summer, fleshy flattened stems and spiny leaves, sandy soil, full sun |
| Cardinal Flower (Lobelia cardinalis) | 3-4' | red tubular flowers, Aug, partial to full sun, prefers moist soil, hummingbirds |
| Clover, Purple Prairie (Dalea purpurea) | 2, | purple, mid-summer, full sun, well drained soil, birds and butterflies |
| Clover, Round-headed Bush (Lespedeza capitata) | 6', | white flowrs, midsummer-fall, dry, well drained full sun, attractive winter feature |
| Columbine, Wild (Aquilegia canadensis) | 1-2' | red and yellow flowers in spring, hummingbirds, well-drained loam, shade or partial sun |
| Compass Plant (Silphium laciniatum) | 7-8' | yellow flowers, July-Sept., huge deeply lobed leaves, tall-grass prairie plant, drought tolerant |
| Coneflower, Pale Purple (Echinacea pallida) | 3-4' | pink flowers, June, drought tolerant, full sun, hummingbirds, birds, butterflies 🛪 🕅 |
| Coneflower, Purple (Echinacea purpurea) | 3-4' | deep pink, June, well-drained loam, high medicinal, full to partial sun, hummers, birds, butterflies |
| Coneflower, Yellow (Ratibida pinnata) | 4-5' | yellow, June-July, well-drained soil, drought tolerant, sun birds, butterflies |
| Coreopsis, Tall (Coreopsis tripteris) | 6-8' | large yellow flowers, Jul-Aug, well-drained soil, drought tolerant, sun, hummingbirds, birds, butterflies |
| Culver's Root (Veronicastrum virginicum) | 4-6' | white showy candelabra-like spires, sharp-toothed, whorles leaves, sun to partial, well drained loam, birds, butter |
| Dock Prairie (Silphium terebinthinaceum) | 6-8' | yellow, heart-shaped basal leaves, Jul-Sept., drought tolerant, full sun, birds, butterflies |
| Golden Alexander (Zizia aurea) | 1-3' | yellow, spring-early summer, full sun, moist to wet soil |
| Golden Ragwort (Senecio aureus) | 1-1.5 | yellow flower, early spring, well-drained soil, partial sun |
| *Goldenrod, Elm-leaved (Solidago ulmifolia) | 3-4' | yellow sprays of flowers in fall, full sun, average to dry soil |
| Goldenrod, Gray (Solidago nemoralis) | 2-3' | yellow, Sept, well-drained soil, sun to partial sun, birds, butterflies |
| Goldenrod, Rigid (Solidago rigida) | 3-4' | yellow clusters of flowers in fall, well-drained soil, drought tolerant, birds, butterflies MA |
| Goldenrod, Rough-leaved (Solidago rugosa) | 2-6' | yellow sprays, September, sun, tolerates a wide range of soils, butterflies, birds M |
| Goldenrod, Showy (Solidago speciosa) | 3-4 | yellow, very showy flower clusters, deep green foliage, sun to partial sun, avg. to well-drained soil 💥 🦜 |
| Goldenrod, Sweeet (Solidago odora) | 2-5' | yellow, anise-scented, plumelike infolorescence, sun to partial sun, dry soil, birds, butterflies |
| Goldenrod, Zig-zag (Solidago flexicaulis) | 1-3' | yellow flower clusters, stem is zigzag, sun to partial sun, average to dry soil, birds, butterflies |
| Illinois Bundleflower (Desmanthus illinoensis) | 2-3' | greenish-white flowers in summer, twirled clusters of seeds in fall, sun, dry soil |
| Indian Physic (Gillenia stipulata) | 2-3' | sm. white to pale pink flowers on ends of branches, May-July, dry or moist soils, shade |
| Indigo, Wild Blue (Baptisia australis) | 3-4' | blue spikes of flowers, May-June, legume, sun, partial shade, moist soil |
| Iris, Blue Flag (Iris versicolor) | 3-4' | showy blue flowers with purple and yellow beards, May-June, sun to partial, wet or moist soils, water gardens |
| Iris, Copper (Iris fulva) | 2-3' | showy copper flowers, June-July, sun to partial sun, moist to wet soil, 🛪 📈 |
| Ironweed (Vernonia altissima) | 4-5' | purple flowers in summer, average, well-drained soil, butterflies |
| Jerusalem Arterchoke (Helianthus tuberosus) | 4-10' | yellow flowers, late summer, leaves rough and hairy, moist soil, full sun, roots edible |
| Joe-Pye Weed (Eupatorium purpureum) | 4-8' | mauve clusters, July-Sept, moist to wet soil, sun to partial sun, butterflies |
| Lobelia, Great Blue (Lobelia siphilitica) | 3-4' | blue tubular flowers, Aug., average to moist soil, full to partial sun |
| Loosestrife, Smooth (Lysimachia quadriflora) | 2-3' | yellow flowers, June-Aug., partial sun, moist soil, sun to partial sun |
| Loosestrife, Wing-angled (Lythrum alatum) | 1-3' | pink flowers at base of leaves, summer, sun to partial shade, wet soils |
| Meadow Rue, Tall (Thalictrum polygamum) | 3-7' | white flowers in billowy masses, late summer, partial shade, well-drained to damp soils |
| Milkweed, Butterfly (Asclepias tuberose) | 1-4' | large orange clusters, June-July, well-drained soil, drought tolerand, clay soils, butterflies, hummers |
| Milkweed, Common (Asclepias syriaca) | 3-5' | rose-purple flowers in large clusters, June-Aug., well-drained soils, full to partial sun |

| Milkweed, Ozark (Asclepias viridis) | 2-3' | light green flowers in clusters, May-July, dry, poor to avg. soil, full to part sun, butterflies, hummers 🛪 🔀 |
|--|----------|--|
| Milkweed, Purple (Asclepias purpurea) | 2-3 | rose to purple flowers, May-Jul, sun to partial sun, moist soil but adapts to garden soil, butterflies, hummers |
| Milkweed, Swamp (Asclepias incarnata) | 3-5' | pink clusters, Jul-Sept, full-pt sun, moist soil, but adapts to garden soil, butterflies, hummers |
| Milkweed, Whorled (Asclepias verticillata) | 1-2' | green, June-Sept, soft-needle-like leaves, avg. garden soil, full sun, butterflies, birds |
| Mint, Hoary Mt. (P. pycnanthemoides) | 1-3' | pink-white, July-Aug, well-drained soil, sun to partial sun |
| Mint, Slender Mountain (P. tenuifolium) | 2-3' | white flowers, summer, sun, average soil, butterflies |
| Mint, Downy Wood (<i>Blephilia ciliata</i>) | 1-1.5' | purple, fuzzy basal leaves, early summer, nice ground cover, well-drained soil, butterflies |
| Mistflower (Eupatorium coelestinum) | 3' | blue-violet, July, average to wet soil, full to partial sun, butterflies |
| Nodding Wild Onion (Allium cernuum) | 1-2' | pink dangling flowers, narrow July-Aug., grass-like leaves, well-drained soil, full to partial sun, butterflies |
| Obedient Plant (Physostegia virginiana) | 3-4' | pink, Aug, moist to dry soil, full sun, birds, butterflies, hummers 🛪 📈 🤼 |
| Petunia, Wild (Ruellia humilis) | 2' | blue-violet, Jul-Aug., full to partial sun, well-drained soil |
| Poppy Mallow, Purple (Callirhoe involucrata) | <u> </u> | fuschia blooms, June-Aug., good full sun ground cover, spreads well, well-drained soil |
| Primrose, Evening (Oenothera biennis) | 6' | light yellow, large flowers, June-Oct., full sun, dry soil |
| Primrose, Missouri (Oenothera missouriense) | 1' | large yellow flowers in summer, good border plant, average soil, butterflies, hummingbirds |
| Primrose, Showy (Oenothera speciosa) | 1' | large pink flowers spring through summer, full sun, good bank stabilizer, butterflies |
| Quinine, Wild (Parthenium integrifolium) | 3-4' | white flower clusters in summer with yucca-like foliage, full sun, drought tolerant, butterflies, hummingbirds |
| Rattlesnake Master (Eryngium yuccifolium) | 3-4' | yucca-like foliage, green flowers in summer, full sun, drought tolerant 🏲 🔀 |
| Rose Mallow (Hibiscus moscheutos) | 4-5' | large funnel-shaped white flowers, mid-summer, full-part sun, avg. to wet soil, hummingbirds, butterflies |
| Rosinweed (Silphium integrifolium) | 3-5' | yellow, July-Sept., well-drained soil, adapted to clay soils, full sun, birds, butterflies |
| Seedbox (Ludwidgia palustris) | 3-4' | small yellow flowers in summer, small seed rattles in fall, full to partial sun, prefers moist soil |
| Senna, Wild (Cassia marilandica) | 3-5' | yellow flowers, late summer, full sun, avg. soil, birds, hummingbirds, butterflies 🛪 📈 |
| Skullcap, Downy (Skutellaria incana) | 2-3' | blue flowers, June-Aug., sun to partial sun, average soil |
| Sneeze-weed, autumn (Helenium autumnale) | 3' | yellow daisy-like petals with yellow centers, Sept-Oct., full sun, mesic to wet soil, butterflies ₩ |
| Spurge, Flowering (Euphorbia corollata) | 2-3' | white, numerous small, resembles baby's breath, July-Aug., avg. to dry, well-drained soil, full sun |
| Spiderwort, Ohio (Tradescantia ohiensis) | 1-2' | violet-blue, April to July, avg to wet soil, full to partial sun |
| Spiderwort, Virginia (Tradescantia virginiana) | 1-2' | violet-blue flowers, Apr-Jul, appear in morning only, shade to full shade, avg. soil |
| Stonecrop (Sedum ternatum) | 4" | star-shaped flowers on fleshy leaves that form low mats, well-drained soil, sun to partial shade, nice groundcover |
| Strawberry, Wild (Fragaria virginiana) | 6" | white flowers, spring, edible berries trailing habit, well-drained soil, sun to partial, good wildlife food |
| Sundrops (Oenothera fruticosa) | 1-2' | lemon yellow flowers all spring, avg. to dry, well-draied soil, drought tolerant, full sun early bloomer 🛪 🕅 |
| Sunflower, Ox-eye (Heliopsis helianthoides) | 3-4' | yellow, June-Sept, full to partial sun, well-drained loam, long blooming period 🗡 📈 |
| Sunflower, Western (Helianthus occidentalis) | 3-4' | many yellow flowers on tall stems above basal leaves, spreads well, full to partial sun, well-drained soil |
| Susan, Sweet (Rudbeckia subtomentosa) | 4' | yellow, July-Sept., sun to partial shade, well drained soil, birds, butterflies 💆 🦫 |
| Thimbleflower (Anemone virginiana) | 3-4' | white flowers, June, well-drained soil, sun to partial sun, butterflies ₩ |
| Turtlehead (Chelone glabra) | 3-4' | white-to pink, Aug-Oct, avg. garden soil, shade to partial shade, butterflies, hummers 🕶 💆 |
| Vervain, Blue (Verbena hastate) | 4-6' | blue-violet spikes of flowers, July, moist to wet soils, full sun |
| Violet, Birdsfoot (<i>Viola pedata</i>) | 6"1 | violet and lavender 2-lipped flowers, leaves shaped like a bird foot, sun to partial sun, well-drained soil |

NATIVE TREES/SHRUBS

| Buttonbush (Cephalanthus occidentalis) | 5-5.5' | multi-stemmed shrub, white spherical flower heads, summer, partial sun, moist to wet soils |
|--|--------|---|
| Hearts Bursting with Love (Euonymous americanus) | shrub | understory shrub with showy drooping pinkish fruits, fall, moist, well-drained organic soil |
| Indigo Bush (Amorpha fruticosa) | 8-12' | shrubby plant with spikes of purple flowers, summer, full sun, moist to wet soil |
| Shrubby St. John's-wort (Hypericum frondosum) | shrub | beautiful yellow flowers mid summer, avg soil, best in full sun, bees, butterflies |
| Trumpetvine (Campsis radicans) | vine | reddish orange trumpet-shaped flowers, July-September, average soil, hummers |

NATIVE WARM SEASON GRASSES ****

| Bottlebrush Grass (Hysterix patula) | 2-4' | long-bristled flower spikelets, June-Aug., shade to partial shade |
|---|------|---|
| Brown Fox Sedge (Carex vulpinoidea) | 2-3' | hardy sedge for moist-wet soils, full to partial sun, small brown and green flowers, birds |
| Buffalo Grass (Buchole dactyloides) | 4-6" | short, fine-textured, heat and cold-resistant, full sun, drought tolerant |
| Dark Green Bulrush (Scirpus atrovirens) | 3-5' | large sedge, rounded dark green-brown flower clusters, moist-wet soils, shallow water |
| Eastern Gamma Grass (Tripsacum dactyloides) | 6' | large coarse grass, sun to partial sun, moist soils |
| June Grass (Koeleria cristata) | 2' | short, slender grass for landscaping in masses, part-full sun, blooms early |
| Path Grass (Juncus tenuis) | 1' | moist to dry, open habitats |
| River Oats (Chasmanthium latifolium) | 2-3' | arching stems with flattened florets, moist soil, partial sun, drought tolerant, nice landscaping plant |
| Sallow Sedge (Carex lurida) | 2-5' | June, moist to wet soils, sun- partial sun, birds 🦥 |
| Virginia Wild Rye (Elymus virginicus) | 2-4' | blue green seed heads, sun-partial shade, sandy to wet soils |

^{****}All of the native warm season grasses provide wonderful food and cover for wildlife and good winter landscaping interest for any yard or garden